

served in Eocene Simsboro deposits in Milam County, Texas.

Annual rainfall of approximately 60 in., channel pattern slightly to highly meandering, average stream gradient of about 5 ft/mi, and bank stabilization by dense vegetation are major parameters controlling Amite deposition of coarse sand and pebble-gravel sediment. Stratification types are related directly to specific depositional features, and include (from thalweg to overbank): thalweg (large-scale cross-stratification), lower point bar (trough cross-stratification, avalanche beds), chute bar (parallel laminae, avalanche beds, trough fill), chute fill (parallel inclined laminae, climbing ripples), and overbank (parallel inclined laminae, mud drape, avalanche beds).

Fundamental differences between point bars of streams transporting coarse-grained bed load and streams having fine-grained bed load are: muddy floodplain deposits are associated only with fine-grained bed-load streams; upper point-bar sediments with ripple cross-stratification and parallel inclined laminae occur only in fine-grained fluvial deposits; chute-front avalanche beds are common in coarse-grained fluvial deposits but are not found in fine-grained fluvial deposits. Coarse-grained fluvial deposits do not show the upward decrease in grain size that is reported to characterize fluvial sediment.

The Simsboro consists mainly of thalweg, lower point-bar, and chute-bar deposits; chute-fill and overbank deposits are preserved only in the last depositional sequence.

D. F. MERRIAM, Kansas Geological Survey, Lawrence, Kan.

INTRODUCTION TO SYMPOSIUM

(No abstract submitted)

A. A. MEYERHOFF, The American Association of Petroleum Geologists, Tulsa, Okla.

CONTINENTAL SHELF POSITIONS DURING GEOLOGIC TIME

Exploration of continental shelves and slopes requires a thorough understanding of the implications of the sea-floor-spreading hypothesis. If the Americas and Eurafica once were joined, as sea-floor spreaders allege, no great intellect is required to see that the geology of the circum-Atlantic shelves and slopes would be different than if the Atlantic Ocean had always existed. Most sea-floor-spreading advocates believe that the Atlantic basin first opened during Jurassic or Early Cretaceous time. If true, drilling for pre-Jurassic objectives in certain areas is senseless. However, if sea-floor spreading has *not* taken place, pre-Jurassic objectives may underlie almost all shelf and slope areas. Similar conclusions can be made for the shelves and slopes surrounding the Indian Ocean.

Sea-floor spreading also would affect the circum-Pacific margins, which should contain strongly deformed post-Jurassic sediments. Drilling objectives in the circum-Pacific shelves should be mainly in young strata, principally of Tertiary age. This seems to be true around most of the Pacific rim, and sea-floor-spreading proponents cite the youthful geology of the circum-Pacific as evidence for their viewpoint.

Although the problem of sea-floor spreading is unusually complex, the problem can be summarized simply: can the hypothesis be proved or disproved? The correct answer to this question will affect world mineral economics for decades to come.

Numerous facts of geology and geophysics contradict the hypothesis of sea-floor spreading. A combination of Ushakov's, Talwani *et al.*'s, and Melson *et al.*'s models of the mid-ocean ridges explains the linear magnetic anomalies of the ridges, their topography, the distribution of rock types on the ridges, their gravity expressions, and the Sykes' "transform-fault" solutions on the basis of *known* processes, as opposed to *inferred* processes. G. D. Afanas'yev's discovery that most metamorphic rocks of the Indian Ocean ridge system have Proterozoic (Riphean) and Paleozoic K-Ar dates severely damages the sea-floor-spreading concept as it has been applied in the Indian Ocean. Rezanov has proved that paleomagnetic methods cannot be used to determine ancient polar positions, or to demonstrate continental separations. Maxwell's, Glangaud *et al.*'s, and Watson and Johnson's studies of the Mediterranean Sea region eliminate both spreading and closing of that sea since late Paleozoic or earlier time. A problem which is even more baffling is that of the *immaculati*: *i.e.*, the presence of flat-lying, undeformed sediments in parts of the deep ocean basins where strong deformation is predicted by sea-floor spreading. Such areas include (1) fracture zones which cross the mid-ocean ridges (undeformed sediments as old as Paleocene are exposed in these fracture zones), (2) abyssal plains, (3) continental slopes and rises (except for gravity-slide blocks), and (4) island-arc trenches. Why does a strongly deformed Early Jurassic to Eocene fold belt extend from the Cape Verde Islands to Tunisia? How does one account for the presence of bathyal to abyssal Jurassic sediments in the Cape Verde Islands, beneath the Hatteras Abyssal Plain, and elsewhere?

Almost conclusive evidence against sea-floor spreading comes from the fields of climatology, meteorology, and biology-paleontology. The field of meteorology is particularly critical, but few geologists, geophysicists, or oceanographers have studied this science. Salomon-Calvi, one of Wegener's staunchest advocates, concluded before his death that widespread continental glaciation would be impossible without an adequate supply of moisture and, therefore, that separation of the southern continents was essential from Mississippian through Permian times. Rukhin's recent analysis of late Paleozoic tillite distribution is even more conclusive because of the wealth of climatology data available to him.

To test Salomon-Calvi's and Rukhin's conclusions, the writer prepared detailed maps of coal, evaporite, and tillite distributions by age and epoch from late Proterozoic time to the present. The evaporite-coal-tillite distribution patterns shown on the maps differ appreciably from previously published compilations, and coincide very closely with the distribution pattern of modern deserts, evaporites, peat bogs, and till deposits. Meteorological and ocean-current analyses of these maps show that the distribution of these rock types *requires*, for the past 1 billion years, an ocean basin-continent distribution nearly the same as that of today. If the continent and ocean-basin distribution of the past was significantly different from that of today, the distribution patterns of pre-Holocene evaporites, coals, and tillites cannot be explained. The maps even show that the position of the horse latitudes has not changed appreciably for 1 billion years. The maps do *not* eliminate some polar tilting since Proterozoic time, but such tilting is not required to explain the patterns.