

2. In nearshore continental beds, species of microfossils are less numerous generally; many beds contain mainly one kind of palynomorph, such as resinous bodies or fusinized material. Possibly this reflects local conditions which favor minimum particle transport and mixing.

3. Bladdered pollen grains occur in varying abundance in almost every bed regardless of other palynomorph abundance. Differing hydrodynamic response of these large grains may govern their anomalous occurrence, in contrast to smaller nonbladdered forms.

DONALD E. OWEN, Dept. Geology, Bowling Green State Univ., Bowling Green, Ohio, and ROY L. INGRAM, Dept. Geology, Univ. North Carolina, Chapel Hill, N.C.

#### PRACTICAL COMPARISON OF METHODS OF COMPUTING GRAIN-SIZE PARAMETERS

Sieved modern coastal sands were used to compute standard grain-size parameters from all known graphic formulas and the method of moments. Progressively greater variation existed in computation of mean, sorting, skewness, and kurtosis. As a group, Folk and Ward's parameters came closest to approximating moment parameters.

The median yielded the greatest deviation from the moment mean, being too fine for coarse-skewed samples and too coarse for fine-skewed samples. Graphic means were very close to the moment mean. Formulas (Folk and Ward, McCammon) considering points near the tails of the distribution and the median produced almost the same values as the moment mean. Formulas (Otto, Inman) omitting the median yielded slightly too coarse means for coarse-skewed samples and slightly too fine for fine-skewed samples. Sampling more than four frequency curve points did not increase accuracy of means or sorting estimates.

Among graphic sorting estimates, Krumbein's  $QD_\phi$  consistently was appreciably lower and Friedman's  $SO_\phi$  and Griffiths'  $PD_\phi$  consistently were appreciably higher than moment standard deviation. Cadigan's  $S$  and Otto's and Inman's  $\sigma_\phi$  produced irregular values averaging higher and lower, respectively, than moment values. Folk and Ward's  $\sigma I$  and McCammon's  $\sigma$  yielded consistent results slightly below moment values. Tanner's  $D_{3,29}$  produced the closest approach to moment values. The non-phi-unit methods (Trask's  $SO$ ; Miller and Zeigler's  $DS$ ; Sharp and Fan's  $Si$ ) showed the same relative change in sorting between samples.

No graphic skewness estimate correlated well with moment skewness, although they usually yielded the same sign. Only Folk and Ward's  $Sk_7$  and Inman's  $\sigma_\phi$  produced fairly consistent results close to the average skewness value. No graphic kurtosis estimate correlated with moment kurtosis, which was consistently higher.

B. PARKASH and G. V. MIDDLETON, Dept. Geology, McMaster Univ., Hamilton, Ont.

#### GRAIN AND GRAPTOLITE ORIENTATION IN TURBIDITE GRAYWACKE, CLORIDORME FORMATION (ORDOVICIAN), GASPÉ, QUEBEC

Sedimentary structures, textures, and fabric were studied in eight graywacke beds exposed on the wave-cut platform near Grande Vallée; these beds were traced continuously for 2 mi along the strike which is almost parallel with the average current direction indicated by sole marks.

The direction of current flow during deposition of

the bed was determined from intrabed lineations and grooves, oriented graptolites, and grain fabric. Most beds show large deviations of internal flow directions from sole-mark directions, particularly in distal regions. Internal directions were increasingly divergent from the sole toward the top of the bed. Agreement between dimensional grain orientations and other internal flow-direction indicators proves that grain orientations are depositional, not tectonic fabrics, and that grains generally were oriented parallel with the flow and imbricated in an upcurrent direction. Current-normal orientation was found in a plane-laminated part of one bed. Grains tend to lack statistically significant preferred orientation in proximal parts of the beds but show preferred orientation as the beds were traced distally.

These results suggest that existing concepts of turbidity-current flow are oversimplified and neglect the possibility of large changes in flow direction, perhaps caused by secondary currents within a single turbidity current.

JAMES M. PARKS, Center for Marine and Environmental Studies, Lehigh Univ., Bethlehem, Pa.

#### MULTIVARIATE FACIES MAPS

Facies maps are constructed from paleontologic and lithologic data to depict major and subtle depositional environment differences across a region during a specified time span. Ratio maps and three-component maps exhibit a lack of discrimination because they cannot incorporate all available data. Factor analysis and cluster analysis techniques can be used to construct truly multivariate facies maps. Earlier attempts at factor or cluster analysis multivariate facies maps had one or more deficiencies: (1) inability to handle a sufficient number of variables and locations; (2) inability to handle mixed-mode data (presence-absence, coded states, integer counts, and continuously variable measurements); (3) inability to take into account redundant or highly correlated variables; and (4) inability to accommodate to missing data.

A new cluster analysis classification computer program has been written to overcome these deficiencies. The FORTRAN IV program can utilize up to 200 variables on as many as 1,000 stations. It performs a distance function principal components analysis to compute orthogonal (uncorrelated) factor measurements for a distance function cluster analysis of locations. This combination will handle mixed-mode data and will adjust to missing data.

From the resulting multivariate classification of paleontologic and lithological data, a facies map showing the distribution of the various classes was constructed and compared with previously published facies maps. An example using multivariate lithologic data from coded AmStrat sample-description logs from central Montana demonstrates the potentialities of this method.

W. T. PECORA, U.S. Geol. Survey, Washington, D.C.  
MINERAL RESOURCE POTENTIAL OF CONTINENTAL MARGIN OF UNITED STATES

The submerged margin of the continent adjacent to the United States is an asset of great importance to the nation. In the aggregate it is a most promising frontier for exploration and development of mineral resources.

The continental margin, considered here to extend

from the coastline to a depth of about 2,500 m, covers an area of about 1.4 million sq mi, approximately a third as large as the land area of the United States. However, knowledge of the geologic framework of this vast area is rudimentary, to say the least.

In broadest terms we know of a great prism of relatively undisturbed sedimentary rocks off the Atlantic and Gulf of Mexico coasts in contrast to a group of narrow basins of sedimentary rocks off the Pacific coast of California and Alaska. The Bering Sea and Arctic coast of Alaska also are underlain by thick accumulations of sedimentary rocks. With a total volume of about 2.5 million cu mi, these sedimentary provinces have a geologic composition, regional geologic framework, and tectonic setting that leave little doubt of a recoverable resource in excess of 100 billion bbl of petroleum liquids and 300 trillion cu ft of gas; and perhaps even in excess of 200 billion bbl of petroleum and 1,000 trillion cu ft of gas. Resources in place are many times greater than these recoverable estimates. Other known resources are salt dome sulfur on the order of several million tons, sand and gravel, shell, phosphorite, and a variety of placer deposits.

To develop these resources to the outer limit of the continental margin for the benefit of the economy of the United States will involve resolution of the worldwide problem of the seaward extent of the jurisdiction of maritime nations. It also will require geologic studies of the offshore areas infinitely more intensive than the studies of the past and the application of continually advancing techniques, both for exploration and ultimate development of the resources. Vision and bold action will be the guidelines.

M. DANE PICARD, Utah Univ., Salt Lake City, Utah, and LEE R. HIGH, JR.,<sup>1</sup> Oberlin College, Oberlin, Ohio

#### LACUSTRINE CRITERIA

Although among the least abundant of depositional environments represented in the geologic record, lakes exceed their limited occurrence in interpretive significance. Lacustrine rocks yield important information concerning paleogeography, paleoclimatology, and tectonics. Therefore geologists should seek to expand their ability to identify and interpret lacustrine environments.

Modern lakes are present in a variety of geologic settings, but most lakes are too small and ephemeral to be geologically significant. Only large, long-lived lakes, such as those in structural basins with interior drainage, can accumulate major lacustrine sedimentary sequences. The recognition of lacustrine rocks requires a variety of techniques. Inasmuch as unique criteria generally are lacking, the geologist must infer origin from scattered, commonly unrelated, and often contradictory data.

Lakes closely resemble shallow, epicontinental seas in physical properties. Large differences, therefore, in lithologic characteristics, sequences, facies relations, sedimentary structures, paleocurrent patterns, and other physical aspects should not be expected. Comparisons of these features from lacustrine and epicontinental rocks indicate the scarcity of significant diagnostic differences.

Large lakes and epicontinental seas differ mainly in size and chemistry. Size differences are evident; few lakes exceed 10,000 sq mi in area. Thus, regional strati-

graphic and paleogeographic relations can be used to differentiate lakes and seas. Geochemical differences, however, are more definitive. Normal seawater has been relatively constant in composition for most of geologic history and changes related to evaporation, precipitation, or dilution are predictable. On the other hand, the chemistry of lacustrine water is not uniform, but is determined partly by lithology and climate in upland source areas. The geochemical balance of lakes, therefore, varies widely in different areas and rarely approximates that of seawater. Accordingly, lakes and seas can differ in authigenic and early diagenetic minerals. Especially useful are evaporite cycles.

Marine and nonmarine environments commonly are distinguished readily by their fauna. Paleontologic differentiation of nonmarine environments is uncertain, however, and requires further study.

ROBERT A. POHOWSKY, Dept. Geology, Univ. Cincinnati, Cincinnati, Ohio

#### STUDIES ON SEDIMENTOLOGY OF SHELL BEDS ON BERMUDIAN PATCH REEFS

The upper surfaces of two patch reefs on the northern edge of the Bermuda platform were observed to consist of coral-coraline algal rock mounds amid a complex system of low-lying channels containing coarse- to fine-grained calcareous detritus. The rock mounds on the larger of the two reefs are noticeably concentrated toward the periphery. Detailed mapping of a part of one of these reefs indicates that the distribution of *Spondylus* valves and other coarse detritus bears a strong relation to the shape and width of the intermound channels. Oddly, 79% of the shells lay with the concave side up. The apparent preference for this generally unstable orientation is attributed to the lofting of shells during transport and the preferential burial of shells lying concave-side-downward.

The mode of the fine fractions of samples taken from shelly areas consistently is toward the right of the mode of samples taken from adjacent nonshelly areas. It thus appears that the shells act as baffles to permit the local deposition of fine sediment.

It is concluded that shell-bed sedimentology reflects the existence of two wave-energy regimes throughout the year. In winter, powerful storm waves determine the distribution of shell beds; calmer seas prevailing in summer months promote the preferential deposition of fines among the shelly material. Further, the rock-sand configuration on larger patch reefs suggests that the reefs may concentrate unimodal sand toward the center and bimodal sediments in the more narrow, circuitous, shell-rich channels toward the periphery.

J. E. RAPSON-McGUGAN, Dept. Geology, Mount Royal Junior College, and Inst. of Sedimentary and Petroleum Geology, Calgary, Alta.

#### INTRA-PERMIAN CORROSION BRECCIAS, SOUTHERN CANADIAN ROCKY MOUNTAINS

An extremely widespread intra-Permian transgression eroded and successively truncated beds from Middle Permian (Guadalupian) to Mississippian (Meramecian) in age.

On this erosion surface are lag gravels in a phosphatic matrix, with associated corrosion breccias. The breccias are developed where planate beds of calcilutite, dololutite, and chert are infiltrated, corroded, and brecciated *in situ* by the phosphate and then partly or totally replaced by chalcedony. Fragments from the breccias may be incorporated in the overlying

<sup>1</sup>Support received through Petroleum Research Fund (Am. Chem. Soc.) Grant (3217-A2).